

Math 20550: Calculus

Name: _____

Practice Final Exam *December 2009*

Section: _____

Record your answers to the multiple choice problems by placing an $\times$ through one letter for each problem on this answer sheet. There are 20 multiple choice questions. Please sign the honor statement if you agree:

"I strictly followed the Notre Dame Honor Code during this test."

Your Signature _____

- | | |
|--|---|
| <p>1. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>2. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>3. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>4. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>5. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>6. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>7. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>8. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>9. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>10. ☐ a ☐ b ☐ c ☐ d ☐ e</p> | <p>11. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>12. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>13. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>14. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>15. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>16. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>17. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>18. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>19. ☐ a ☐ b ☐ c ☐ d ☐ e</p> <p>20. ☐ a ☐ b ☐ c ☐ d ☐ e</p> |
|--|---|

Record your answers to the multiple choice problems by placing an $\times$ through one letter for each problem on this answer sheet. There are 20 multiple choice questions. Please sign the honor statement if you agree:

"I strictly followed the Notre Dame Honor Code during this test."

Your Signature _____

1. ☐ a ☐ b ☐ c ☐ d ☐ e

11. ☐ a ☐ b ☐ c ☐ d ☐ e

2. ☐ a ☐ b ☐ c ☐ d ☐ e

12. ☐ a ☐ b ☐ c ☐ d ☐ e

3. ☐ a ☐ b ☐ c ☐ d ☐ e

13. ☐ a ☐ b ☐ c ☐ d ☐ e

4. ☐ a ☐ b ☐ c ☐ d ☐ e

14. ☐ a ☐ b ☐ c ☐ d ☐ e

5. ☐ a ☐ b ☐ c ☐ d ☐ e

15. ☐ a ☐ b ☐ c ☐ d ☐ e

6. ☐ a ☐ b ☐ c ☐ d ☐ e

16. ☐ a ☐ b ☐ c ☐ d ☐ e

7. ☐ a ☐ b ☐ c ☐ d ☐ e

17. ☐ a ☐ b ☐ c ☐ d ☐ e

8. ☐ a ☐ b ☐ c ☐ d ☐ e

18. ☐ a ☐ b ☐ c ☐ d ☐ e

9. ☐ a ☐ b ☐ c ☐ d ☐ e

19. ☐ a ☐ b ☐ c ☐ d ☐ e

10. ☐ a ☐ b ☐ c ☐ d ☐ e

20. ☐ a ☐ b ☐ c ☐ d ☐ e

1. Let S be the part of cylinder $y^2 + z^2 = 1$, with $z \geq 0$, and $0 \leq x \leq 1$, and let S have the upward orientation. Determine which of the following equals $\int \int_S \mathbf{F} \cdot d\mathbf{S}$ where $\mathbf{F}(x, y, z) = \langle 0, 0, z \rangle$.

(a) $\int_0^1 \int_{-1}^1 \sqrt{1-y^2} dy dx.$

(b) $\int_0^1 \int_{-1}^1 \sqrt{1-x^2} dy dx$

(c) $\int_0^1 \int_{-1}^1 (1-y^2) dy dx$

(d) $\int_0^1 \int_{-1}^1 [\sqrt{1-y^2}]^{-1} dy dx$

(e) $\int_0^1 \int_{-1}^1 (1-x^2) dy dx$

2. Find the maximum rate of change of $f(x, y) = x^2y + 2y$ at the point $(-1, 2)$ and the direction in which it occurs.

(a) The maximum rate of change is 5 in the direction $\frac{1}{5}\langle -4, 3 \rangle$.

(b) The maximum rate of change is 5 in the direction $\frac{1}{5}\langle -3, 4 \rangle$.

(c) The maximum rate of change is 5 in the direction $\frac{1}{5}\langle 4, 3 \rangle$.

(d) The maximum rate of change is 5 in the direction $\frac{1}{5}\langle 4, -3 \rangle$.

(e) The maximum rate of change is 6 in the direction $\frac{1}{5}\langle -4, 3 \rangle$.

3. Evaluate $\int_C (1 + x^2y) ds$ where C is the upper half of the unit circle $x^2 + y^2 = 1$.

(a) $\pi + \frac{2}{3}.$

(b) $2\pi.$

(c) $0.$

(d) $\frac{2}{3}.$

(e) $1.$

4. Determine which of the following integrals gives the volume of the region bounded by the cylinder $x^2 + y^2 = 1$, and the planes $z = 0$ and $x + z = 1$.

(a) $\int_0^{2\pi} \int_0^1 \int_0^{1-r \cos \theta} r dz dr d\theta$ (b) $\int_0^\pi \int_0^1 \int_0^{1-r \cos \theta} r dz dr d\theta$
 (c) $\int_0^{2\pi} \int_0^1 \int_0^{1-r \cos \theta} dz dr d\theta$ (d) $\int_0^{\frac{\pi}{2}} \int_0^1 \int_0^{1-r \cos \theta} r dz dr d\theta$
 (e) $\int_0^1 \int_0^{\sqrt{1-y^2}} \int_0^{1-x} dz dx dy$

5. Let C be the curve $\mathbf{r}(t) = \langle t, \cos(2t), 1 + \sin(3t) \rangle$, $0 \leq t \leq \frac{\pi}{2}$, and let

$$\mathbf{F}(x, y, z) = \langle y(2x + z), x(x + z) - z, y(x - 1) + 2z \rangle.$$

Compute $\int_C \mathbf{F} \cdot d\mathbf{r}$. (Hint: Find f with $\mathbf{F} = \nabla f$).

(a) $-\frac{\pi^2}{4}$ (b) 0 (c) $\frac{\pi}{2}$ (d) $1 - \frac{\pi}{4}$ (e) $\frac{\pi^2}{2} - 1$

6. If $z = f(x, y)$, $x = u^2 + v^2$ and $y = u^2 - v^2$, find $\frac{\partial^2 z}{\partial u \partial v}$.

(a) $4uv(f_{xx} - f_{yy})$ (b) $2(f_{xx} + f_{yy})$ (c) $4uv(f_{xx} + f_{yy})$
 (d) $4uv(f_{xx} + 2f_{yy} - f_{yy})$ (e) $4uv(f_{xx} + 2f_{yy} + f_{yy})$

7. Find the scalar projection, $\text{comp}_{\mathbf{v}}(\mathbf{w})$, of the vector $\mathbf{w} = \langle 1, 1, 2 \rangle$ onto the vector $\mathbf{v} = \langle 2, -2, 1 \rangle$.

(a) $\frac{2}{3}$ (b) $-\frac{2}{3}$ (c) 1 (d) 2 (e) $\frac{2}{\sqrt{6}}$

8. Determine which of the following integrals gives the area of the region in the xy -plane below the x -axis above $y = x^2 - 2$ and to the left of $y = -2x - 2$.

(a) $\int_{-2}^0 \int_{-\sqrt{y+2}}^{-\frac{y}{2}-1} dx dy$ (b) $\int_{-2}^0 \int_{-\sqrt{y+2}}^{1-\frac{y}{2}} dx dy$ (c) $\int_{-2}^0 \int_{\sqrt{y+2}}^{-\frac{y}{2}-1} dx dy$
 (d) $\int_{-\sqrt{2}}^0 \int_{-\sqrt{y+2}}^{-\frac{y}{2}-1} dx dy$ (e) $\int_{-2}^0 \int_{-2x-2}^{x^2-2} dy dx$

9. Find surface area of the part of the paraboloid $z = x^2 + y^2$ that lies under the plane $z = 4$.

(a) $\int_0^{2\pi} \int_0^2 r\sqrt{1+4r^2}drd\theta$

(b) $\int_0^{2\pi} \int_0^2 \sqrt{1+4r^2}drd\theta$

(c) $\int_0^\pi \int_0^2 r\sqrt{1+4r^2}drd\theta$

(d) $\int_0^{2\pi} \int_0^1 r\sqrt{1+4r^2}drd\theta$

(e) $\int_0^{2\pi} \int_0^4 r\sqrt{1+4r^2}drd\theta$

10. Let C be the triangle with vertices $(0, 0)$, $(1, 1)$, and $(2, 0)$ oriented counterclockwise. Compute

$$\int_C [\cos x^{100} + x^4 y^5]dx + [\sin(e^y) + x^5 y^4]dy$$

- (a) 0 (b) 1 (c) -1 (d) $\frac{4}{5}$ (e) $\frac{5}{4}$

11. Express the area between $x^2 + \frac{y^2}{9} = 1$ and $x^2 + \frac{y^2}{9} = 9$ as an integral, using the substitution $x = r \cos \theta$ and $y = 3r \sin \theta$.

(a) $\int_0^{2\pi} \int_1^3 3rdrd\theta$

(b) $\int_0^{2\pi} \int_1^3 9rdrd\theta$

(c) $\int_0^{2\pi} \int_1^3 3r^2drd\theta$

(d) $\int_0^{2\pi} \int_1^9 3rdrd\theta$

(e) $\int_0^{2\pi} \int_0^3 3rdrd\theta$

12. Calculate the arc length of the helix parameterized by $\mathbf{r} = \langle -3t, 4 \cos t, -4 \sin t \rangle$ for $0 \leq t \leq \pi$

- (a) 5π (b) 0 (c) 10π (d) 12π (e) 2π

13. Find the absolute minimum of $f(x, y) = x^2 + 2y^2 + 4y - 2$ on the disk $x^2 + y^2 \leq 4$.

- (a) -4 (b) -6 (c) 0 (d) -2 (e) 14

14. Find a direction vector for the line of intersection of the planes $x + y + 2z = 1$ and $3x - y = 0$.

- (a) $\langle 1, 3, -2 \rangle$ (b) $\langle 1, 3, 2 \rangle$ (c) $\langle 3, -2, 1 \rangle$ (d) $\langle 3, 1, -2 \rangle$ (e) $\langle -2, 3, 1 \rangle$

15. Let C be the intersection of the cylinder $x^2 + y^2 = 1$ and the plane $z = 2y + 3$ oriented counter-clockwise with the normal upwards. Use Stokes Theorem to calculate $\int_C \mathbf{F} \cdot d\mathbf{r}$ where

$$\mathbf{F}(x, y, z) = \langle 2e^y - z, \cos(yz), xe^y \rangle.$$

- (a) 2π (b) $\sqrt{5}\pi$ (c) $\frac{\pi}{\sqrt{5}}$ (d) $\frac{2\pi}{\sqrt{5}}$ (e) 0

16. Evaluate $\int \int \int_E 3e^{[(x^2+y^2+z^2)^{\frac{3}{2}}]} dV$ where E is the upper *half* of the ball radius 1 centered at the origin.

- (a) $2\pi(e-1)$ (b) $\pi(e-1)$ (c) $4\pi(e-1)$ (d) π (e) 2π

17. Find the minimum of the function $f(x, y, z) = x^2 + y^2 + 2z^2$ on the surface $x^2y^2z = 16$.

- (a) 10 (b) 8 (c) 16 (d) 12 (e) 14

18. A particle starts from rest at time $t = 0$ at the origin $(0, 0, 0)$. It then begins to move with acceleration $\mathbf{a}(t) = \langle 1, 6t, 12t^2 \rangle$. Find the time, if ever, at which the particle passes through the point $(1, 1, 1)$.

- (a) never (b) $t = 1$ (c) $t = 2$ (d) $t = 3$ (e) $t = 6$

19. Determine two vectors that are tangent to the surface $\mathbf{r}(u, v) = \langle vu^2 - 2u, uv^2 - v, uv \rangle$ at the point $(0, 1, 2)$.

- (a) $\langle 2, 1, 1 \rangle, \langle 4, 3, 2 \rangle$ (b) $\langle 2, 4, 2 \rangle, \langle 1, 3, 1 \rangle$ (c) $\langle 0, 1, 1 \rangle, \langle 4, 1, 1 \rangle$

- (d) $\langle 1, 2, -1 \rangle, \langle 1, -2, 1 \rangle$ (e) $\langle 0, 2, -1 \rangle, \langle 1, -6, 3 \rangle$

20. Find $\int \int_S \mathbf{F} \cdot d\mathbf{S}$ where $\mathbf{F}(x, y, z) = \langle xy, \frac{3}{4}y, -zy \rangle$ and S is the surface of the sphere $x^2 + y^2 + z^2 = 4$ with the outward orientation.

- (a) 8π (b) π (c) 2π (d) 16π (e) 0